

GMRT Observations of Low z Damped $\text{Ly}\alpha$ Absorbers

Jayaram N. Chengalur^{1*} and Nissim Kanekar^{1†}

¹*National Center for Radio Astrophysics (TIFR), University of Pune Campus, P. O. Bag 3, Ganeshkhind, Pune 411007, India.*

ABSTRACT

We present Giant Metrewave Radio Telescope (GMRT) observations of redshifted HI 21cm absorption in two low redshift ($z = 0.2212$, $z = 0.0912$) damped $\text{Ly}\alpha$ systems seen towards the Gigahertz peaked source OI 363 ($z_{em} = 0.630$). The object at $z = 0.0912$ is the lowest redshift damped $\text{Ly}\alpha$ system known to date. Ground based imaging (Rao & Turnshek 1998) shows that at neither redshift is there a large spiral galaxy at low impact parameter to the line of sight to OI 363, in contradiction with the suggestion that damped $\text{Ly}\alpha$ systems are large proto-disks. Since OI 363 is a highly compact, core dominated source, the covering factor of the HI gas is likely to be unity. Nonetheless, the spin temperatures derived from the 21cm optical depth (and using the N_{HI} measured from HST spectra (Rao & Turnshek 1998)) are high, viz. 1120 ± 200 K and 825 ± 110 K for the high and low redshift systems respectively. These values are considerably higher than typical values (100 – 200 K) measured in our Galaxy and Andromeda and are, in fact, similar to those obtained in high redshift damped $\text{Ly}\alpha$ systems. Our observations hence suggest that evolutionary effects may not be crucial in understanding the difference in derived spin temperature values between local spiral disks and high redshift damped $\text{Ly}\alpha$ systems.

Key words: quasars: absorption lines, galaxies: evolution, ISM: general, cosmology: observations

1 INTRODUCTION

Damped $\text{Ly}\alpha$ systems are the extremely high HI column density ($N_{HI} > 2 \times 10^{20}$ atoms cm^{-2}) systems seen in absorption in the spectra taken towards distant quasars. Although rare, they are the dominant contributors (by mass) to the observed neutral gas at high ($z \sim 3$) redshifts. Principally for this reason, these systems are natural candidates for the precursors of $z = 0$ galaxies. Consistent with this interpretation, the mass density of neutral gas in damped $\text{Ly}\alpha$ systems at $z \sim 3$ is comparable to the mass density in stars in luminous galaxies at $z = 0$. Thus, to zeroth order, the evolution with redshift of the neutral gas density matches that expected from gas depletion due to star formation (eg. Lanzetta et al. 1991, Lanzetta, Wolfe & Turnshek 1995, Storrie-Lombardi, McMahon & Irwin 1996). Further, the evolution of metallicity with redshift also roughly matches what one would

expect from models of galactic evolution (Ferrini, Mollá & Díaz 1997, Fall 1997).

On the other hand, the morphology of damped $\text{Ly}\alpha$ systems remains poorly understood. Based on the edge-leading asymmetries seen in the absorption profiles of low ionization metals associated with these systems, Prochaska & Wolfe (1997) suggest that they are rapidly rotating large disks with significant vertical scale heights. However, such profiles can also be explained by models in which damped systems are much smaller objects that are undergoing infall and merger (Haehnelt, Steinmetz & Rauch 1997). It has also been claimed that the metal abundance of damped $\text{Ly}\alpha$ systems depends on the total HI column density in a way as would be expected from large disks with central HI holes (Wolfe & Prochaska 1998), although the number of systems involved in this study is small.

For damped $\text{Ly}\alpha$ systems that lie in front of radio loud quasars, it is possible to augment the optical/UV spectra with HI 21cm absorption spectra. Such a comparison, yields, among other things (and under suitable assumptions), the spin temperature, T_s , of the HI gas. Derived spin tempera-

* chengalur@ncra.tifr.res.in

† nissim@ncra.tifr.res.in

tures of damped Ly α systems have, in general, been much larger than those observed in the disk of the Galaxy or in nearby galaxies (Braun & Walterbos 1992, Braun 1996), implying that either damped Ly α systems are not disks, or that the ISM in the damped Ly α proto-disks is considerably different from that in the local $z = 0$ disks, presumably due to evolutionary effects.

Studies of low redshift damped Ly α systems are particularly interesting in this regard, since evolutionary effects are expected to be negligible. Further, much more detailed information is obtainable, in particular from HST and/or ground based imaging, which makes identification of the absorber possible (eg. Le Brun et al. 1997, Lanzetta et al. 1997). Of course, it remains a possibility that the population that gives rise to damped Ly α absorption at low redshift is distinct from that at high redshift.

In this paper, we report the detection of redshifted 21cm absorption in two low redshift ($z = 0.2212$, $z = 0.0912$) damped Ly α systems seen towards the quasar OI 363 (0738+313, $z_{em} = 0.630$) (Rao & Turnshek 1998). Observations of the higher redshift system confirm, at considerably improved spectral resolution and sensitivity, earlier results from the Westerbork Synthesis Radio Telescope (WSRT) (Lane et al. 1998), while the lower redshift ($z = 0.0912$) system is the lowest redshift damped Ly α system known to date.

2 OBSERVATIONS AND DATA REDUCTION

The observations were carried out using the GMRT (Swarup et al. 1991, Swarup et al. 1997). The backend used was the proto-type eight station FX correlator, which gives a fixed number (128) of spectral channels over a total bandwidth that can be varied from 64 kHz to 16 MHz. Due to various ongoing maintenance and installation activities, the actual number of antennas that were available during our observing runs varied between six and eight.

For the observations of the $z = 0.0912$ system the bandwidth was set to 1.0 MHz. No spectral taper was applied, giving a channel spacing of $\sim 1.8 \text{ km s}^{-1}$. Two observing runs were made, one on 27 June 1998 and the other on 5 July 1998. The on source time for each run was about six hours. Two observing runs were also taken for the $z = 0.2212$ system (on 26 June 1998 and 4 July 1998), the first with a total bandwidth of 1.0 MHz (i.e. a channel spacing of $\sim 2.0 \text{ km s}^{-1}$) and the other with a total bandwidth of 0.5 MHz (i.e. a channel spacing of $\sim 1.0 \text{ km s}^{-1}$). Each of these observing runs had an on source time of ~ 4 hours. Bandpass calibration at both redshifts was done using 3C 295, which was observed at least once during each observing run.

The data was converted from the raw telescope format to FITS and then reduced in AIPS in the standard way. Maps were produced after subtracting out the continuum emission of the background quasar using UVLIN, and spectra extracted from the resulting three dimensional cube. The GMRT does not do online doppler tracking; this is, however, unimportant since the doppler shift within any one of our observing runs was a small fraction of a channel. For the




Figure 1. GMRT redshifted 21cm absorption spectrum of the lower redshift system towards OI363. The channel spacing is $\sim 1.8 \text{ km s}^{-1}$. The deepest optical depth (~ 0.18) is at a heliocentric redshift of 0.09118. The width (FWHM) of the line is $\sim 5 \text{ km s}^{-1}$.

lower redshift system, data from the observations on different days were corrected to the heliocentric frame and then combined.

The final spectrum for the $z = 0.0912$ system is shown in Figure 1. There appear to be two components, one considerably deeper than the other. The fainter component, although weak, was detected in both our observing runs, and its magnitude is also considerably higher than the noise level. It is, of course, possible that the spectrum consists of two components, one of which is broad and weak and the other, much deeper but narrow. The redshift of the narrow component is consistent with the redshift quoted in Rao & Turnshek (1998). The peak optical depth is ~ 0.18 (i.e. a depth of 390 mJy with the continuum flux of OI363 being 2.0 Jy), and occurs at a redshift of $z = 0.09118 \pm 0.00001$. The FWHM of the line is small, $\sim 5 \text{ km s}^{-1}$.

Lane et al. (1998) report a redshift of $z = 0.2212$ for the higher redshift system, based on WSRT observations. The 2.0 km s^{-1} GMRT spectrum (which has a considerably better velocity resolution and sensitivity than the WSRT spectrum) is shown in Figure 2. The redshift measured from this spectrum is 0.2212 ± 0.00001 . This is consistent with the redshift measured from the 1.0 km s^{-1} resolution spectrum (which is not shown here). The peak optical depth (~ 0.07) is somewhat less than that of the lower redshift system, but the velocity width is comparable, $\sim 5.5 \text{ km s}^{-1}$ (FWHM).




Figure 2. GMRT redshifted 21cm absorption spectrum of the higher redshift system towards OI363. The channel spacing is $\sim 2.0 \text{ km s}^{-1}$. The peak optical depth (~ 0.07) occurs at a heliocentric redshift of 0.2212. The width (FWHM) of the line is $\sim 5.5 \text{ km s}^{-1}$

3 DISCUSSION

The total HI column density of a damped system can be determined from its Ly α profile; this can be then used, in combination with the measured HI 21cm optical depth, to determine the spin temperature of the absorbing gas (under the assumption that the absorbing gas is homogeneous. For a multi-phase absorber, this derived temperature is the column density weighted harmonic mean of the spin temperatures of the different phases, provided all the phases are optically thin.). One of the principal uncertainties in this derivation of the spin temperature is the covering factor of the absorbing gas. In particular, the radio emission from quasars is often extended, while the UV continuum source is essentially a point source; thus, the line of sight along which the HI column density has been derived from UV measurements need not be the same as the one for which the 21cm optical depth has been measured. The case of OI 363, however, is relatively straightforward. OI 363 is a core dominated Gigahertz peaked source whose total flux decreases from 2.2 Jy at 1.64 GHz to 1.59 Jy at 408 MHz. VLA maps at 1.64 GHz (Murphy, Browne & Perley 1993) and 1.4 GHz (Saikia et al. 1998) show that the source is highly core dominated, with about 97% of the total flux in an unresolved core component. VLBI measurements at 5 GHz (Stanghellini et al. 1996) show that the core size is ~ 10 milli arcseconds ($\sim 16 \text{ pc}$ and $\sim 31 \text{ pc}$ at redshifts of 0.0912 and 0.2212 respectively, for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$). Stanghellini et al. (1997) note that their 5 GHz VLBI map

recovers only 77% of the total flux measured by the VLA at 5 GHz. While it is unclear whether there is much change in the source size between 5 GHz and 1.3 GHz, we know from IPS measurements (Jeykumar et al. 1998) that the upper limit on the core size at 330 MHz is 50 milli arcseconds. At both redshifts, the depth of the line (see Figures 1 & 2) considerably exceeds the flux in the weak lobes, this implies that the absorbers must cover the central core. Given the small size of this central core, the covering factor is likely to be close to unity.

The HI column densities inferred from the present observations, in terms of the spin temperatures of the two damped systems, are $1.82 \pm 0.02 \times 10^{18} T_s \text{ atoms cm}^{-2}$ and $0.71 \pm 0.04 \times 10^{18} T_s \text{ atoms cm}^{-2}$, for the lower and higher redshift systems, respectively. The column densities measured by Rao & Turnshek (1998), from the damped Lyman- α lines, are $7.9 \pm 1.4 \times 10^{20} \text{ atoms cm}^{-2}$ and $1.5 \pm 0.2 \times 10^{21} \text{ atoms cm}^{-2}$, again in order of decreasing redshift. The spin temperatures obtained are hence $825 \pm 110 \text{ K}$ (for the $z = 0.0912$ absorber) and $1120 \pm 200 \text{ K}$ (for the $z = 0.2212$ system). For the higher redshift system, our measurement agrees within the errors with that of Lane et al. (1998). The overwhelming source of the (formal) uncertainty is in the determination of the HI column density from the UV measurements. Thus, even at redshifts where no evolution is expected, the derived spin temperature is significantly higher than that typically seen in the Galaxy. If one assumes that the HI 21cm spectral width is entirely due to thermal motions, the required kinetic temperatures are $\sim 625 \text{ K}$ and $\sim 750 \text{ K}$ for the lower and higher redshifted system respectively, i.e. comparable to the derived spin temperatures. Note however, that in the ISM of the Galaxy, there is no stable neutral phase with temperature $\sim 1000 \text{ K}$. On the other hand, such high spin temperatures appear common at both high and intermediate redshifts (see eg. de Bruyn, O’Dea & Baum 1996, Carilli et al. 1996, Lane et al. 1996, Kanekar & Chengalur 1997, Boissé et al. 1998).

Ground based imaging of the OI 363 field (Rao & Turnshek 1998) shows that there are no spiral galaxies at small impact parameters to the line of sight, contrary to the canonical model where damped systems arise in extended disks. Similarly, the next lowest redshift damped Ly α absorber (0850+4400, Lanzetta et al. 1997) appears to be associated with an S0 galaxy, while, at intermediate redshifts, the absorbers appear to be associated with galaxies spanning a wide range of morphological types (Le Brun et al. 1997). Interestingly, at lower redshifts still, where imaging of HI 21cm emission is possible, 21cm absorption from quasar galaxy pairs appears to be associated more with tidal tails or other extended features of gas rich galaxies (Carilli & van Gorkom 1992), and not directly with the disks of large spirals.

While the low number density of damped Lyman- α systems at $z < 1$ makes it *a priori* extremely unlikely that two such systems might be found along the same line of sight, the VLBI map of OI 363 appears to rule out the possibility of this line of sight being biased due to gravitational lensing. The current observations (and the absence of detectable gravitational lensing) do not however place strong

constraints on the surface density or mass of the absorbing systems.

In summary, it appears that even at the lowest redshifts, gas outside the disks of spiral galaxies and with apparent physical parameters considerably different from the ISM of nearby galaxies has a non-trivial contribution to the total absorption cross-section. This is consistent with observations that, even for intermediate redshift damped Ly α absorbers, the metallicity is considerably lower than typical solar values (Boissé et al. 1998). Finally, the present GMRT observations also suggest that evolutionary effects may not play an important role in understanding why the derived spin temperature for damped Ly α systems are in general higher than those measured in nearby spiral galaxy disks.

Acknowledgments These observations would not have been possible without the many years of dedicated effort put in by the GMRT staff to build the telescope. The GMRT 1400 MHz wide-band feed and receiver system was built by the Raman Research Institute. We are also grateful to Wendy Lane, Judith Irwin, Anish Rosh, D. J. Saikia, R. Srianand and Kandu Subramanian for their comments and suggestions.

REFERENCES

- Braun R., 1997, ApJ, 484, 637
 Braun R., Walterbos R., 1992, ApJ, 386, 120
 Boissé P., Le Brun V., Bergeron J., Deharveng J-M., 1998, A&A, submitted
 Carilli C. L., van Gorkom J. H., 1992, ApJ 399, 384
 Carilli C. L., Lane W., de Bruyn A. G., Braun R., Miley G. K., 1996, AJ, 111, 1830
 de Bruyn A. G., O’Dea C. P., Baum, S. A., 1996, A&A 305, 450
 Fall S. M., 1997, Petitjean P., Charlot S., eds., Structure and Evolution of the Intergalactic Medium from QSO Absorption Line Systems, Editions Frontières
 Ferrini F., Mollá M., Díaz A. I., 1997, ApJ, 487, L29
 Haehnelt M. G., Steinmetz M., Rauch M., 1998, ApJ, submitted
 Jeykumar S. J., Saikia D. J., Rao A. P., Balasubramanian, V., 1998, in preparation
 Kanekar N., Chengalur J. N., 1997, MNRAS, 292, 831
 Lane W., Smette A., Briggs F., Rao S., Turnshek D., Meylan G., 1988, ApJ, in press
 Lanzetta, K.M., Wolfe, A.M., Turnshek, D.A. Lu, L., McMahon, R.G. & Hazard, C., 1991, ApJS, 77,1
 Lanzetta K. M., Wolfe A. M., Turnshek D. A., 1995, ApJ, 440, 435
 Lanzetta K. M., Wolfe A. M., Altan H., Barcons X., Chen H-W., Fernández-Soto A., Meyer D. M., Ortiz-Gil A., Savaglio S., Webb J. K. & Yahata N., 1997, AJ, 114, 1337
 Le Brun V., Bergeron J., Boissé P., Deharveng J-M., 1997, A&A, 321, 733
 Murphy D. W., Browne I. W. A., Perley R. A., 1993, MNRAS, 264, 298
 Prochaska J. X., Wolfe A. M., 1997, ApJ, 487, 73
 Rao S., Turnshek D. A., 1998, ApJL, in press
 Saikia D. J., Holmes G. F. Kulkarni A. R., Salter C. J., Garrington S. T. 1998, MNRAS, in press
 Stanghellini, C., Dallacasa D., O’ Dea C. P., Baum S. A., Fanti R., Fanti C., 1997, A&A, 325, 943
 Storrie-Lombardi L. J., McMahon R. G., Irwin M. J., 1996, MNRAS, 283, L79
 Swarup G., Ananthakrishnan S., Kapahi V. K., Rao A. P., Subrahmanya C. R., Kulkarni V. K. 1991, Current Science, 60, 95
 Swarup G., Ananthakrishnan S., Subrahmanya C. R., Rao A. P., Kulkarni V. K., Kapahi V. K. 1997, Jackson N., Davis R. J. eds. in High-sensitivity Radio Astronomy, Cambridge
 Wolfe A. M., Prochaska J. X., 1998, ApJ in press

Figure 1. GMRT redshifted 21cm absorption spectrum of the lower redshift system towards OI363. The channel spacing is $\sim 1.8 \text{ km s}^{-1}$. The deepest optical depth (~ 0.18) is at a heliocentric redshift of 0.09118. The width (FWHM) of the line is $\sim 5 \text{ km s}^{-1}$.

Figure 2. GMRT redshifted 21cm absorption spectrum of the higher redshift system towards OI363. The channel spacing is $\sim 2.0 \text{ km s}^{-1}$. The peak optical depth (~ 0.07) occurs at a heliocentric redshift of 0.2212. The width (FWHM) of the line is $\sim 5.5 \text{ km s}^{-1}$.

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.